

**EVALUATION OF ANTIDIABETIC ACTIVITY OF ALCOHOLIC
EXTRACTS OF *HALDINA CORDIFOLIA* LEAF ON
STREPTOZOTOCIN INDUCED DIABETIC RATS**

**A Dissertation submitted to
The Tamil Nadu Dr.M.G.R Medical University Chennai**

In partial fulfillment of the degree of

MASTER OF PHARMACY

(Pharmacology)

Submitted by

SIJAD T.K

(Reg No :261526155)

Under the guidance of

Prof.G. NAGARAJA PERUMAL, M.Pharm.,

**Professor & Head,
Department of Pharmacology
Karpagam College of Pharmacy
Coimbatore-32**



MAY 2017

**DEPARTMENT OF PHARMACOLOGY
KARPAGAM COLLEGE OF PHARMACY
COIMBATORE-641 032**

CERTIFICATES

CERTIFICATE

This is to certify that this dissertation entitled “***evaluation of antidiabetic activity of alcoholic extracts of haldina cordifolia leaf on streptozotocin induced diabetic rats***” Submitted by **Mr. Sijad T.K** to The Tamil Nadu Dr.M.G.R Medical University, Chennai in partial fulfilment for the degree of **Master Of Pharmacy In Pharmacology** is a bonafied work carried out by the candidate under my guidance and supervision in the Department of Pharmacology, Karpagam College of Pharmacy Coimbatore-32.

I have fully satisfied with his performance and work. I have forward this dissertation work for evaluation.

Station:
Date :

Mr.G.NAGARAJA PERUMAL M.Pharm.,
Professor & Head
Department of Pharmacology

CERTIFICATE

This is to certify that this dissertation entitled “***evaluation of antidiabetic activity of alcoholic extracts of haldina cordifolia leaf on streptozotocin induced diabetic rats***” Submitted by **Mr. SIJAD T.K** to The Tamil Nadu Dr.M.G.R Medical University, Chennai in partial fulfilment for the degree of **Master Of Pharmacy In Pharmacology** is a bonafied work carried out under the guidance and supervision of **Mr.G.Nagaraja Perumal,M.Pharm.**, Professor & Head, Department of Pharmacology, Karpagam College of Pharmacy Coimbatore-32.

I have fully satisfied with his performance and work. I have forward this dissertation work for evaluation.

Station:

Dr. S. MOHAN, M.Pharm., Ph.D.,

Date:

Principal

DECLARATION

I hereby declare that this dissertation entitled “***evaluation of antidiabetic activity of alcoholic extracts of haldina cordifolia leaf on streptozotocin induced diabetic rats***” submitted by me, in partial fulfilment of the requirements for the degree of **Master Of Pharmacy In Pharmacology** to The Tamil Nadu Dr.M.G.R Medical university, Chennai is the result of my original and independent research work carried out under the guidance of **Mr.G. Nagaraja Perumal, M.Pharm.**, Professor & Head Department of Pharmacology, Karpagam College of Pharmacy, Coimbatore-32, & Co-Guide **Dr. R.S. Parihar, Banner PharmacapS**, during the academic year 2016- 2017.

Station:

Mr. SIJAD T.K

Date:

EVALUATION CERTIFICATE

This is to certify that the dissertation work entitled “***evaluation of antidiabetic activity of alcoholic extracts of haldina cordifolia leaf on streptozotocin induced diabetic rats***” submitted by, **Mr. SIJAD T.K** bearing Reg.No: **261526155** to The Tamil Nadu Dr. M. G. R Medical University, Chennai in partial fulfilment for the Degree of **Master Of Pharmacy In Pharmacology** is a bonafied work carried out during the academic year 2013-2014 by the candidate at Department of Pharmacology, Karpagam College of Pharmacy, Coimbatore and was evaluated by us.

Examination centre:

Date:

Internal Examiner

Convener of Examination

External examiner

**DEDICATED TO MY BELOVED PARENTS,
TEACHERS AND ALMIGHTY**

ACKNOWLEDGEMENT

ACKNOWLEDGEMENT

First of all I would like to thank god for his blessings to do this research work successfully. With immense pleasure and pride, I would like to take this opportunity in expressing my deep sense of gratitude to my beloved guide **Mr G. Nagaraja Perumal M.Pharm.**, Professor & Head, Department of Pharmacology, Karpagam College of Pharmacy under whose active guidance, innovative ideas, constant inspiration and encouragement of the work entitled “***evaluation of antidiabetic activity of alcoholic extracts of haldina cordifolia leaf on streptozotocin induced diabetic rats***” has been carried out.

I take this opportunity with pride and immense pleasure expressing my deep sense of gratitude to my respectable co-guide **Dr R.S. Parihar**, General Manager, of BANNER PHARMACAPS, whose innovative ideas, active guidance, inspiration, tremendous efforts, encouragement, help and continuous supervision has made the dissertation a grand and glaring success to complete.

I wish to express my deep sense of gratitude to Dr.R.Vasanthakumar Chairman of Karpagam Group of institutions for the facilities provided me in this institution.

My sincere thanks to our respected and beloved Principal Dr.S.Mohan, M Pharm ,Ph.D, Karpagam College of Pharmacy for his encouragement and also providing all facilities in this institution to the fullest possible extent enabling me to complete this work successfully.

I convey my gratitude to Mrs.V.Idachristi M. Pharm,Professor & Head ,Department of Pharmacognosy helped me to proceed useful ideas.

My whole hearted thanks to Mr.D. Ranjith kumar M Pharm,Asst. Professor,Department of Pharmaceutical Analysis for his kind advice.

I am also conveying my thanks to Mrs. M. Karpagavalli ,M. Pharm, Associate Professor, Department of Pharmaceutical chemistry, for encouragement and valuable suggestion during this work.

I express my sincere thanks to Mr. K. Nahas , Lab assistant , Department of Pharmaceutical chemistry for his kind support.

I convey my gratitude to Mr. S. Asker , Lab Assistant , Department of Pathology for his kind support.

I express my sincere thanks to Mrs.M. Sathybhama Lab assistant, Department of Pharmaceutical chemistry for her kind support.

I am duly bound to all my non teaching staffs of Karpagam collage of Pharmacy for their valuable advices and co-operation. Above all , I am remain indebted to my seniors class mates (Anoop, Bhavan,Shanavas,Amritha,Habeeb,mohammed shanavas v.k, Ubaid), to my beloved parents who inspired and guided me and also for being tha back bone for all my successful endeavors in my life.

SIJAD.TK

(261526155)

CONTENTS

CH.NO	CONTENTS	PAGE NO.
1	INTRODUCTION	1
2	REVIEW OF LITERATURE	9
3	AIM AND OBJECTIVE	14
4	PLAN OF WORK	15
5	PLANT PROFILE	16
6	MATERIALS AND METHODS	21
7	RESULTS AND DISCUSSION	31
8	SUMMARY AND CONCLUSION	45
9	BIBLIOGRAPHY	48

LIST OF TABLE

SL NO.	TITLE	PAGE NO.
1	α-Amylase Inhibition of Petroleum ether extract of <i>Haldina cordifolia</i> leaf	31
2	α-Amylase Inhibition of Ethyl acetate extract of <i>Haldina cordifolia</i> leaf	32
3	α-Amylase Inhibition of Ethanolic extract of <i>Haldina cordifolia</i> leaf	33
4	α-Amylase Inhibition of Acarbose (Possitive control)	34
5	TLC results of ethanolic extract of <i>Haldina cordifolia</i> leaf	35
6	Preliminary phytochemical screening of ethanolic extract of <i>Haldina cordifolia</i> leaf	37
7	Hypoglycemic Test	39
8	Results of the effects of ethanolic extract on blood Glucose levels	41
9	Oral Glucose Tolerance Test	43

LIST OF FIGURES

FIG NO.	TITLE	PAGE NO.
1	Main Symptoms of Diabetes	2
2	Plant of <i>Haldina Cordifolia-I</i>	16
3	Plant of <i>Haldina Cordifolia-II</i>	16
4	Plant of <i>Haldina Cordifolia-III</i>	17
5	Plant of <i>Haldina Cordifolia-IV</i>	17
6	Plant of <i>Haldina Cordifolia-V</i>	18
7	Plant of <i>Haldina Cordifolia-VI</i>	18
8	Plant of <i>Haldina Cordifolia-VII</i>	19
9	αAmylase Inhibition of Petroleum ether extract of <i>Haldina cordifolia</i> leaf	32
10	αAmylase Inhibition of Ethyl acetate extract of <i>Haldina cordifolia</i> leaf	33
11	αAmylase Inhibition of Ethanolic extract of <i>Haldina cordifolia</i> leaf	34
12	αAmylase Inhibition of Acarbose (Possitive control)	35

CHAPTER-I

1. INTRODUCTION

1.1 Diabetes Mellitus (DM)

The Diabetes Mellitus is chronic metabolic disorder, which there are high blood sugar level over a prolonged period. The term "mellitus" or "from honey" was added by the Briton John Rolle in the late 1700s to separate the condition from diabetes insipidus, which is also associated with frequent urination. It is a fast-growing global problem with huge social, health and economic consequences. It is estimated that in 2010 there were globally 285 millions people (approximately 6.4% of the adult population) suffering from this disease. This number is estimated to increase to 430 million in the absence of better control or cure. An ageing population and obesity are two main reasons for the increase. Furthermore it has been shown that almost 50% of the putative diabetics are not diagnosed until 10 years after onset of the disease, hence the real prevalence of global diabetes must be astronomically high.

This high blood sugar produces the symptoms of frequent urination, increased thirst, and increased hunger. Untreated, diabetes can cause many complications. Acute complications include diabetic ketoacidosis and non ketotic hyperosmolarcoma. Serious long-term complications include heart disease, stroke, kidney failure, foot ulcers and damage to the eyes.

Indian physicians around the same time identified the disease and classified it as Madhumeha or "honey urine", noting but the urine would attract ants. This is possibly due to the diet and life-style of the ancient people, or because the clinical symptoms were observed during the advanced stage of the disease. Galen named the disease "Diarrhea of the urine" (diarrhea urinosa). The earliest surviving work with a detailed reference to diabetes is that of Aretaeus of Cappadocia (2nd or early 3rd century CE). Type 1 and type 2 diabetes were identified as separate conditions for the first time by the Indian physicians Sushruta and Charaka in 400-500 CE with type 1 associated with youth and type 2 with being overweight. Effective treatment was not developed until the early part of the 20th century, when Canadians Frederick

Banting and Charles Herbert Best isolated and purified insulin in 1921 and 1922. This was followed by the development of the long-acting insulin NPH in the 1940s.

1.2 Signs and Symptoms

Fig No:1- Main Symptoms of Diabetes



1.3 Overview of the most significant symptoms of diabetes

The classic symptoms of untreated diabetes are weight loss, polyuria (frequent urination), polydipsia (increased thirst), and polyphagia (increased hunger). Symptoms may develop rapidly (weeks or months) in type 1 diabetes, while they usually develop much more slowly and may be subtle or absent in type 2 diabetes.

Several other signs and symptoms can mark the onset of diabetes, although they are not specific to the disease. In addition to the known ones above, they include blurry vision, headache, fatigue, slow healing of cuts, and itchy skin. Prolonged high blood glucose can cause glucose absorption in the lens of the eye, which leads to changes in its shape, resulting in vision changes. A number of skin rashes that can occur in diabetes are collectively known as diabetic dermadromes.

People (usually with type 1 diabetes) may also experience episodes of diabetic ketoacidosis, a type of metabolic problems characterized by nausea, vomiting and abdominal pain, the smell of acetone on the breath, deep breathing known as Kussmaul breathing, and in severe cases a decreased level of consciousness.

A rare but equally severe possibility is hyperosmolar nonketotic state, which is more common in type 2 diabetes and is mainly the result of dehydration.

1.4 Complications

All forms of diabetes increase the risk of long-term complications. These typically develop after many years (10–20), but may be the first symptom in those who have otherwise not received a diagnosis before that time.

The major long-term complications relate to damage to blood vessels. Diabetes doubles the risk of cardiovascular disease and about 75% of deaths in diabetics are due to coronary artery disease. Other "macrovascular" diseases are stroke, and peripheral vascular disease.

The primary microvascular complications of diabetes include damage to the eyes, kidneys, and nerves. Damage to the eyes, known as diabetic retinopathy, is caused by damage to the blood vessels in the retina of the eye, and can result in gradual vision loss and potentially blindness. Damage to the kidneys, known as diabetic nephropathy, can lead to tissue scarring, urine protein loss, and eventually chronic kidney disease, sometimes requiring dialysis or kidney transplant. Damage to the nerves of the body, known as diabetic neuropathy, is the most common complication of diabetes. The symptoms can include numbness, tingling, pain, and altered pain sensation, which can lead to damage to the skin. Diabetes-related foot problems (such as diabetic foot ulcers) may occur, and can be difficult to treat, occasionally requiring amputation. Additionally, proximal diabetic neuropathy causes painful muscle wasting and weakness.

There is a link between cognitive deficit and diabetes. Compared to those without diabetes, those with the disease have a 1.2 to 1.5-fold greater rate of decline in cognitive function.

1.5 Diagnosis

Diabetes mellitus is characterized by recurrent or persistent hyperglycemia, and is diagnosed by demonstrating any one of the following:

- Fasting plasma glucose level ≥ 7.0 mmol/l (126 mg/dl)
- Plasma glucose ≥ 11.1 mmol/l (200 mg/dl) two hours after a 75 g oral glucose load as in a glucose tolerance test
- Symptoms of hyperglycemia and casual plasma glucose ≥ 11.1 mmol/l (200 mg/dl)
- Glycated hemoglobin (Hb A1C) $\geq 6.5\%$.

A positive result, in the absence of unequivocal hyperglycemia, should be confirmed by a repeat of any of the above methods on a different day. It is preferable to measure a fasting glucose level because of the ease of measurement and the considerable time commitment of formal glucose tolerance testing, which takes two hours to complete and offers no prognostic advantage over the fasting test. According to the current definition, two fasting glucose measurements above 126 mg/dl (7.0 mmol/l) is considered diagnostic for diabetes mellitus.

Per the World Health Organization people with fasting glucose levels from 6.1 to 6.9 mmol/l (110 to 125 mg/dl) are considered to have impaired fasting glucose. People with plasma glucose at or above 7.8 mmol/l (140 mg/dl), but not over 11.1 mmol/l (200 mg/dl), two hours after a 75 g oral glucose load are considered to have impaired glucose tolerance. Of these two prediabetic states, the latter in particular is a major risk factor for progression to full-blown diabetes mellitus, as well as cardiovascular disease. The American Diabetes Association since 2003 uses a slightly different range for impaired fasting glucose of 5.6 to 6.9 mmol/l (100 to 125 mg/dl).

Glycated hemoglobin is better than fasting glucose for determining risks of cardiovascular disease and death from any cause.

The rare disease diabetes insipidus has similar symptoms to diabetes mellitus, but without disturbances in the sugar metabolism (*insipidus* means "without taste" in Latin) and does not involve the same disease mechanisms.

Table No .1

WHO DIABETES DIAGNOSTIC CRITERIA			
Condition	2 hour glucose	Fasting glucose	HbA_{1c}
Unit	mmol/l(mg/dl)	mmol/l(mg/dl)	%
Normal	<7.8 (<140)	<6.1 (<110)	<6.0
Impaired fasting glycaemia	<7.8 (<140)	≥ 6.1(≥110) & <7.0(<126)	6.0–6.4
Impaired glucose tolerance	≥7.8 (≥140)	<7.0 (<126)	6.0–6.4
Diabetes mellitus	≥11.1 (≥200)	≥7.0 (≥126)	≥6.5

1.6 Prevention and Treatment

There is no known preventive measure for type 1 diabetes. Type 2 diabetes can often be prevented by a person being a normal body weight, physical exercise, and

following a healthy diet. Dietary changes known to be effective in helping to prevent diabetes include a diet rich in whole grains and fiber, and choosing good fats, such as polyunsaturated fats found in nuts, vegetable oils, and fish. Limiting sugary beverages and eating less red meat and other sources of saturated fat can also help in the prevention of diabetes. Active smoking is also associated with an increased risk of diabetes, so smoking cessation can be an important preventive measure as well.

Diabetes mellitus is a chronic disease, for which there is no known cure except in very specific situations. Management concentrates on keeping blood sugar levels as close to normal ("euglycemia") as possible, without causing hypoglycemia. This can usually be accomplished with diet, exercise, and use of appropriate medications (insulin in the case of type 1 diabetes; oral medications, as well as possibly insulin, in type 2 diabetes).

Learning about the disease and actively participating in the treatment is vital for people with diabetes, since the complications of diabetes are far less common and less severe in people who have well-managed blood sugar levels. The goal of treatment is an HbA1C level of 6.5%, but should not be lower than that, and may be set higher. Attention is also paid to other health problems that may accelerate the deleterious effects of diabetes. These include smoking, elevated cholesterol levels, obesity, high blood pressure, and lack of regular exercise. Specialised footwear is widely used to reduce the risk of ulceration, or re-ulceration, in at-risk diabetic feet.

1.7 Lifestyle

People with diabetes can benefit from education about the disease and treatment, good nutrition to achieve a normal body weight, and sensible exercise, with the goal of keeping both short-term and long-term blood glucose levels within acceptable bounds. In addition, given the associated higher risks of cardiovascular disease, lifestyle modifications are recommended to control blood pressure.

1.8 Medications

Metformin is generally recommended as a first line treatment for type 2 diabetes, as there is good evidence that it decreases mortality. Routine use of aspirin, however, has not been found to improve outcomes in uncomplicated diabetes. Angiotensin

converting enzyme inhibitors (ACEIs) improve outcomes in those with DM while the similar medications angiotensin receptor blockers (ARBs) do not.

Type 1 diabetes is typically treated with a combinations of regular and NPH insulin, or synthetic insulin analogs. When insulin is used in type 2 diabetes, a long-acting formulation is usually added initially, while continuing oral medications. Doses of insulin are then increased to effect.

In those with diabetes some recommend blood pressure levels below 120/80 mmHg; however, evidence only supports less than or equal to somewhere between 140/90 mmHg to 160/100 mmHg.

1.9 Pancreatic transplantation

A pancreas transplant is occasionally considered for people with type 1 diabetes who have severe complications of their disease, including end stage renal disease requiring kidney transplantation.

1.10 Support

In countries using a general practitioner system, such as the United Kingdom, care may take place mainly outside hospitals, with hospital-based specialist care used only in case of complications, difficult blood sugar control, or research projects. In other circumstances, general practitioners and specialists share care in a team approach. Home telehealth support can be an effective management technique.

1.11 Diets And Exercise Treatment

Making healthy food choices is very important to help keep your blood glucose level under control. People with diabetes don't need to buy or prepare special foods. The foods that are best for someone with diabetes are excellent choices for everyone: foods that are low in fat, salt, and sugar, and high in fiber, such as beans, fruits, vegetables, and whole grains. These foods help you reach and stay at a weight that's good for your body. Regular physical activity is important for people with diabetes. Being physically active has been shown to improve blood glucose levels in older people whose levels are high. Exercise is especially good for people with diabetes because it helps control weight, helps insulin work better to lower blood glucose, is good for your heart and lungs, gives you more energy, Regular physical activity improves insulin resistance and lipid profile (reduction in

triglyceride and increase in high-density lipoprotein (HDL)) and lowers blood pressure (although blood pressure will rise during exercise), the metabolic benefits in type 2 diabetes are lost within 3-10 days of stopping regular exercise, physical activity also protects against the development of type 2 diabetes.

1.12 A1C Test

The A1C test is used to detect type 2 diabetes and prediabetes but is not recommended for diagnosis of type 1 diabetes or gestational diabetes. The A1C test is a blood test that reflects the average of a person's blood glucose levels over the past 3 months and does not show daily fluctuations. The A1C test is more convenient for patients than the traditional glucose tests because it does not require fasting and can be performed at any time of the day. The A1C test result is reported as a percentage. The higher the percentage, the higher a person's blood glucose levels have been. A normal A1C level is below 5.7 percent. An A1C of 5.7 to 6.4 percent indicates prediabetes. People diagnosed with prediabetes may be retested in 1 year. People with an A1C below 5.7 percent may still be at risk for diabetes, depending on the presence of other characteristics that put them at risk, also known as risk factors. People with an A1C above 6.0 percent should be considered at very high risk of developing diabetes. A level of 6.5 percent or above means a person has diabetes.

1.13 Herbal drugs

Allopathic medicines are very costly. In contrast, herbal medicines are very cheap. This cost effectiveness makes them all the more alluring. Herbal medicines can be brought without prescription and they are available in almost all health stores. Some herbs can even be grown at home. For certain ailments, herbal medicines are considered to be more effective than allopathic medicines.

Herbal medicines do not have any side effects, as they are free from chemicals. They are also milder than allopathic medicines. The natural detoxification process of the body is effectively enhanced by herbal medicines. They can be used to cleanse the colon, improve digestion and food absorption. Herbal medicines are also very good in boosting the immune system.

Herbal medicines are very effective in curing various digestive disorders like colitis, indigestion, peptic ulcers, and irregular bowel movements. These types of medicines are best for people who are allergic to various types of drugs. Herbal medicines are also effective in boosting the mental health. Most of the ailments related to blood circulation like high blood pressure, varicose ulcers, and many others can be controlled through herbal medicine. Some herbal medicines are very good in reducing the cholesterol level in the blood stream. They are also used to treat coronary artery diseases. Herbal medicine can be used to reduce weight by regulating appetite.

CHAPTER-II

2. REVIEW OF LITERATURE

Ososki et al., (2002)^[22] Reported Ethnobotanical literature survey of medicinal plants in the Dominican republic used for women's health condition on the usage of *Haldina Cordifolia* decotion For uro-genital problems can use as decotion preparation of seeds taken orally or wash bath. The field work in the Dominican republic revealed that herbs were used to treat many women's health condition including uterine fibrosis, excessive uterine bleeding, endometriosis and hot flushes in menopause.

Dasgupta et al., (2013)^[23] investigated and reported The present on Preliminary phytochemical studies of *Haldina Cordifolia* about the important phytochemicals present in the leaf extract. The preliminary screening showed the presence of many phytochemicals such as alkaloids, flavonoids, terpenoids, saponin, fixed oils and fats. High Performance Thin Layer Chromatography analysis was carried out with optimized solvent system consisting of ethyl acetate, formic acid, glacial acetic acid and water in the ratio of 8:1:1:2. The densitometric scanning of the chromatograms of hydroalcoholic extract showed 7 peaks at 254 nm 8 peaks at 366 nm. The phytochemicals detected in the present study justify the therapeutic uses of the leaves in the traditional medicines.

Isaac et al., (2007)^[24] conducted histopathological and biochemical studies on Short-term administration of an aqueous extract of *Haldina Cordifolia Integra* var. *crenata* (Andr.) in a 14-day sub-acute toxicity studies using female Sprague-Dawley rats on *Haldina Cordifolia Integra* (Ki) leaf extract for the

treatment of ulcers, pain and adenoma of the prostate gland reported about its traditionally use for orally administered multipurpose in Ghana and other parts of the world.

Cruz et al., (2012) ^[28] evaluated and studied on the topic [Phytomedicine: International Journal of Phytotherapy & Phytopharmacology](#) about Aqueous extract of *Haldina Cordifolia Pinnata* (Kp) have been found effective in models to reduce acute anaphylactic reactions. They investigated on the effect of Kp and the flavonoid quercetin (QE) and quercitrin (QI) on mast cell activation *in vitro* and in a model of allergic airway disease *in vivo*. Treatment with Kp and QE *in vitro* inhibited degranulation and cytokine production of bone marrow-derived mast cells following IgE/FcεRI crosslinking, whereas treatment with QI had no effect. Similarly, *in vivo* treatment with Kp and QE decreased development of airway hyperresponsiveness, airway inflammation, goblet cell metaplasia and production of IL-5, IL-13 and TNF. In contrast, treatment with QI had no effect on these parameters. These findings demonstrate that treatment with Kp or QE is effective in treatment of allergic airway disease, providing new insights to the immunomodulatory functions of this plant. *Haldina Cordifolia Pinnata* inhibits mast cell activation and prevents allergic airway disease.

Shazid et al., (2012) ^[30] reported on the elucidated structure under the title Chemical and biological studies of *Haldina Cordifolia Pinnata* (Lam.) growing in Bangladesh. In this study they isolated compounds from *K. Pinnata* and elucidated their structures and to explore preliminary antioxidant, antimicrobial, cytotoxic and thrombolytic activities of extractives of the plant. The methanol extract of whole plant of *K. Pinnata* has been subjected to different chromatographic separation and purification processes to isolate the secondary metabolites. The structures of the isolated compounds have been elucidated by extensive NMR studies.

Patil et al., (2013) ^[31] reported under Antidiabetic activity of *Haldina Cordifolia Pinnata* in streptozotocin-induced diabetic rats by glucose independent insulin secretagogue action about the anti diabetic action of *K. Pinnata* in which it was reported that it reduces fasting blood glucose level

Gad et al., (2006) ^[33] studied on Biochemical study of the anti-diabetic action of the Egyptian plants fenugreek and balanites. It proceeded by evaluating the effects of 21 days oral administration of Fenugreek seed and Balanites fruit

extracts (1.5 g/kg bw) on the liver and kidney glycogen content and on some key liver enzymes of carbohydrate metabolism in STZ-diabetic rats were studied. In addition, the effects of these two plant extracts on the intestinal alpha-amylase activity in vitro and starch digestion and absorption in vivo were also examined. .

Patil et al., (2013) ^[41] reported on hypoglycemic activity on *Haldina Cordifolia Pinnata* Lam. (Crassulaceae) under the title Antidiabetic activity of *Haldina Cordifolia Pinnata* in streptozotocin-induced diabetic rats by glucose independent insulin secretagogue action. The probable mechanism for insulin secretagogue action was evaluated through studies using diazoxide and nifedipine. The bioactive component from DCM fraction was studied using HPTLC, GCMS and IR.

Dzhafarova et al., (2009) ^[28] reported on Antidiabetic action of extract of *Juglans Regia* L. The study is on the impact of *Juglans Regia* L on the pathological processes in alloxan-induced diabetic animals. The investigation showed that extract from walnut leaves decreases the blood sugar level, has a positive impact on lipid metabolism. Antioxidant properties of an extract from leaves of the walnut tree (*Juglans regia*) are reported.

Radhika et al., (2011) ^[36] Did a major evaluation on Antidiabetic and Hypolipidemic Activity of *Punica Granatum* Linn on Alloxan Induced Rats the study aimed to evaluate the antidiabetic and hypoglycemic activity of *Punica granatum*. Diabetes & hyperlipidemia was induced by the intra peritoneal injection of alloxan monohydrate (120mg/kg) for 2 consecutive days. Diabetes was confirmed 2 days after the last alloxan dose administration by determining the blood glucose concentration. Treatments were started after confirmation of diabetes in rats. During diabetes, the excess glucose present in the blood reacts with hemoglobin to form glycosylated hemoglobin. So the total hemoglobin level is lowered in alloxan induced diabetic rats. Alloxan induced diabetes has been observed to cause a massive reduction of the beta cells of the islets of pancreas leading to hyperglycemia. Rats treated with alloxan (120mg/kg), for 2 consecutive days, showed an increase in the concentration of glucose, triglycerides, cholesterol, LDL cholesterol, VLDL cholesterol and a decrease in the level of HDL cholesterol and hemoglobin content. Administration of crude powder of *Punica granatum* husk reduced the concentration of glucose, triglycerides, cholesterol, LDL cholesterol, VLDL cholesterol and raised

the level of HDL cholesterol and hemoglobin content in the blood of both group-I normal and group III alloxan diabetic rats treated.

Saif-Ur-Rehman et al., (2011)^[38] presented the work on Antidiabetic Effect of Aloe vera Extract on Alloxan Induced Diabetic Rats. It was carried out to investigate the antidiabetic effect of Aloe vera extract in normal and alloxan induced diabetic rats.

Kuroda et al., (2005)^[40] studied on the turmeric (*Curcuma Longa L.* rhizomes) EtOH extract the Hypoglycemic effects of turmeric (*Curcuma Longa L.* rhizomes) on genetically diabetic KK-Ay mice it shows significant suppression and an increase in blood glucose level in type 2 diabetic KK-A(y) mice.

Mondal et al., (2008)^[27] did a preliminary and antidiabetic screening on the title Antidiabetic activity of *Diospyros Malabarica* Kostel bark: a preliminary investigation for possible mode of action The extract exerted significant antihyperglycemic effect in alloxan-induced hyperglycemia and resulted in increase in plasma protein content and decrease in alkaline phosphatase, cholesterol and triglyceride levels when compared with those in the diabetic control group. However there were no significant changes in body and kidney weights of the DM extract-treated animals, compared to those of the untreated diabetic rats as a control.

Grover et al., (2002)^[40] conducted a research of Medicinal plants of India with anti-diabetic potential on the plants mentioned in ancient literature or used traditionally for diabetes have shown anti-diabetic property. The present paper reviews 45 such plants and their products (active, natural principles and crude extracts) that have been mentioned/used in the Indian traditional system of medicine and have shown experimental or clinical anti-diabetic activity.

Rajnish Gupta et al., (2011)^[46] reported on Antidiabetic efficacy of *Mangifera Indica* seed kernels in rats. The study was conducted to examine the hypoglycemic potency of seed kernels of *Mangifera indica* ethanol extract (MI EtE) in streptozotocin diabetic rats. Remarkable abnormalities were observed in serum and tissue parameters in hyperglycemic rats after streptozotocin administration

Sabjan Khaleel Basha and Vinoji Sugantha Kumari.(2012)^[50] reported on In vitro antidiabetic activity of *Psidium Guajava* seeds extracts. In this they did the

evaluation of the glucose uptake of (antidiabetic activity) crude n-hexane, ethanol, ethanol and aqueous leaf extracts of *P.guajava*. Methods: *P.guajava* leaf extracts were subjected to inhibitory effect of glucose utilization using specific standard in vitro procedure.

Yoshino et al., (2009)^[42] reported on the title Anti-diabetic activity of a leaf extract prepared from *Salacia Reticulata* in mice The effects of a water extract prepared from the leaves of *Salacia reticulata* on the absorption of sugars in normal and type 1 diabetic mice were investigated.

Akhani et al., (2004)^[22] reported on antidiabetic study of *Zingiber officinale* Roscoe as Anti-diabetic activity of *Zingiber Officinale* in streptozotocin-induced type I diabetic rats The fresh and dried rhizome of *Zingiber officinale* Roscoe (commonly known as ginger) is widely used in traditional medicine.The study is on the effect of the juice of *Z. officinale* (4 mL kg(-1), p.o. daily) for 6 weeks on streptozotocin (STZ)-induced type I diabetic rats with particular reference to the involvement of serotonin (5-hydroxytryptamine; 5-HT) receptors in glycaemic control.

Ramadan et al., (2013)^[28] observed two parameters on the diabetic induced rat on title Hypoglycemic and hepatoprotective activity of *Rosmarinus Officinalis* extract in diabetic rats The present study examined the effect of water extract (200 mg/kg body weight) of *Rosmarinus officinalis* L. in streptozotocin (STZ)-induced diabetic rats for 21 days. The hepatoprotective effects were investigated in the liver tissues sections.

CHAPTER-III

3. AIM AND OBJECTIVE

The Diabetes Mellitus is worldwide problem due to the lifestyle changes lead to decreased physical activity, increased consumption of fat, sugar and calories, and higher stress levels, affecting insulin sensitivity and obesity. Based on the above causative factor to increased tenfold, from 1.2% to 12.1%, between 1971 and 2000 entire world. It estimates is estimated that 61.3 millions people aged 20-79 years live with diabetes in India (2015). This number is expected to increase to 101.2 millions by 2030. There are many synthetic drugs used for the treatment of diabetes such as Glibenclamide-sulphonylureas in India and combination therapy because its more side effects reported. Several literatures indicated that the herbal drugs are less adverse effect when compared with synthetic drugs. *Haldina Cordifolia* one of most traditionally using hypoglycemic drug among tribes in india and its same species *K.Pinnata* was scientifically reported but *Haldina Cordifolia* is not scientifically validated. Based on the above mentioned reasons, one new research is required to develop one new drug with activity and less side effects, which will be work multifactorial antidiabetic mechanism. Allopathic medicines are very costly contrast to herbal medicines are very cheap. This cost effectiveness makes them all the more alluring. The work provides scientific validation for the use of leaf against diabetes by revealing the chemical compounds may be present in the plant. The prediction of biological activity of the chemical compounds present in the extract wills supports the *invitro* results after its phytochemical analysis. Herbal drugs were expelled milde side

effects reported in literatures. The present study is attempt to develop a novel plant based antidiabetic drug, which will be evaluated by using *Invitro* and *Invivo* methods

CHAPTER-IV

4. PLAN OF WORK

1. Collection and authentication of the genuine plant material.
2. Shade drying and granulation of the dried leaves.
3. Sequential extraction of the plant material using petroleum ether, ethyl acetate and ethanol.
4. *Invitro* antidiabetic study using α -amylase inhibition assay.
5. Selection of active extract with higher activity.
6. Phytochemical analysis of active extract.
7. Acute toxicity study.
8. *Invivo* antidiabetic study of active extract.
 - I) Normal animal
 - a) OGTT
 - b) Hypoglycemic
 - II) Streptazocin induced diabetes mellitus
 - a) Multiple dose study

CHAPTER-V

5.PLANT PROFILE



Fig No 2. Plant of *Haldina Cordifolia-I*



Figure No 3. Plant of *Haldina Cordifolia*-II



Fig No. 4. Plant of *Haldina Cordifolia*-III



Fig No. 5. Plant of *Haldina Cordifolia*-IV



Fig No. 6. Plant of *Haldina Cordifolia*-V



Fig No. 7. Plant of *Haldina Cordifolia*-VI



Fig No. 8. Plant of *Haldina Cordifolia*-VII

5.1 *Haldina Cordifolia*

Herbalism or "herbal medicines" have a long history to cure several kinds of human diseases from the various parts of the plants such as leaf, stem, bark, leaf, etc. [2] Plants have been the basis for medical treatments through much of human history, and such traditional medicine is still widely practiced today. Modern medicine recognizes herbalism as a form of alternative medicine, as the practice of herbalism is not strictly based on evidence gathered using the scientific method. Many of the pharmaceuticals currently available to physicians have a long history of use as herbal remedies, including opium, aspirin, digitalis, and quinine. The World Health Organization (WHO) estimates that 80 percent of the population of some Asian and African countries presently uses herbal medicines for some aspect of primary health care. The use of, and search for, drugs and dietary supplements derived from plants have accelerated in recent years. Pharmacologists, microbiologists, botanists, and natural-products chemists are combining the Earth for phytochemicals and leads that could be developed for treatment of various diseases. Among the 120 active compounds currently isolated from the higher plants and widely used in modern medicine today, 80 percent show a positive correlation between their modern therapeutic use and the traditional use of the plants from which they are derived. More than two thirds of the world's plant species at least 35,000 of which are estimated to have medicinal value come from the developing countries [1] . It was realized from the various reports that *Haldina cordifolia*, belongs to the family Rubiaceae, have displayed plethoras of potential biological activities such as anticancer, hepatoprotective, antiinflammatory, antifertility, antidiabetic, antiamoebic, antinociptive etc. from its various parts . Furthermore, this plant have also used in curing various ailments such as rheumatism, stomachache, headache, cold/cough, toothache, fever, pain and swelling, bacterial infection, urinary problems, conjunctivitis, miscarriage etc. Keeping the view of the above mentioned medicinal importance of this plant, we therefore became interested to investigate the antibacterial activity of this plant. To the best of our knowledge, the antibacterial activity from this plant has not been reported so far till now. In the present paper, we would like to explore the antibacterial activity of the methanolic extract this plant employing various kinds of strains. It was found that majority of these strains employed, have displayed potential antibacterial activity. Introduction to the plant *Adina cordifolia* Roxb. (Synonym: *Haldina cordifolia*) Family: Rubiaceae Vernacular names: English : Yellow teak 1 Sanskrit : Haridru Hindi : Haldu Telugu :

Pasupukadamba Kannada: Hethega, Arisina tega Morphology A hulking deciduous tree with the shoot often irregularly fluted and buttress at the bottom and normally reaching 6m girth and 15-35m in height. It is a blossoming plant in the family Rubiaceae. Bark to a degree dark grey, thick, bumpy or blackish and to a certain extent pale ashy or silvery white, wood yellow moderately rigid and even-grained. Leaves opposite 10-25cm width, orbicular, abruptly acuminate, pedestal cordate, glabrous above pubescent beneath, with 5-8 pairs of sideways nerves. Petiole 3.8-10cm extended stout, juvenile. Stipules 1.3-1.8cm long oblong or obovate. Flowers is irrelevant individually but is very attractive when they blossom together in balls with a perimeter of 2 to 3cm. They are usually yellow in shade often tinged with a shade of pink. Fruits capsules split into two dehiscent cocci. Seeds many slender small tailed above.

CHAPTER-VI

MATERIALS AND METHODS

6.1.1 Plant collection and authentication

The leafs of *Haldina Cordifolia* were collected from Nilgiri hills Gudalur, Kerala and which was authenticated by Dr.Raghu.A.V Scientist-B,Extension & Training, Kerala Forest Research institute Peechi – 680 653, Thrissur.

6.1.2 Preparation of coarse powder and Extraction technique

The leafs were shade dried at room temperature for 10 days. Then these were milled into powder by mechanical grinder. This powder was sequentially extracted to their increasing polarity with Petroleum ether, Ethyl acetate, Ethanol respectively. About 500gm of powdered leaf was uniformly packed into a thimble in a soxhlet apparatus and extracted with 1000ml Petroleum ether, Ethyl acetate and Ethanol, respectively. Constant heat was provided by Mantox heater for recycling of the solvent. The process of extraction continues for 1-2 hours for each solvent. The

excess solvent was evaporated and the dried extracts were kept in refrigerator at 4°C for their future use in phytochemical analysis and pharmacological screenings.

6.1.3 *Invitro* antidiabetic activity of *Haldina Cordifolia* leaf extracts

6.2 Alpha-amylase inhibition assay

6.2.1 Chemicals or reagents

Potato starch, trichloroacetic acid, Folin-Ciocalteu reagents were purchased from SD Fine Pvt. Ltd., Mumbai, 3,5-dinitrosalicylic acid, Tris buffer, linoleic acid, ammonium molybdate, were purchased from Hi-Media Pvt. Ltd., Mumbai, α -amylase, α -glucosidase enzymes, xanthine oxidase, quercetin, hypoxanthine, pyrocatechol were purchased from SRL Pvt. Ltd., Mumbai. Glucose assay kit from Agappe diagnostic Pvt. Ltd., Kerala, Acarbose was obtained from Bicon Pvt. Ltd., Chennai, ferrozine, (2'2'-azobis (2-amidino propane) dihydrochloride), butylated hydroxy toluene from Loba Cheme. All other chemicals used in the study were obtained commercially and were of analytical grade.

6.2.2 Instrument used

UV-visible Spectrometer (Systronic double beam- UV-2201).

6.2.3 Preparation of extract

Leaf extraction used in *invitro* and *invivo* studies were prepared by using suitable solvents (Carboxy methyl cellulose).

6.2.4 Experimental procedure for α -amylase inhibition assay

A total of 500 μ l of test samples and standard drug (100-1000 μ g/ml) were added to 500 μ l of 0.20 mM phosphate buffer (pH 6.9) containing α -amylase (0.5mg/ml) solution and were incubated at 25°C for 10 min. After these, 500 μ l of a 1% starch solution in 0.02 M sodium phosphate buffer (pH 6.9) was added to each tube. The reaction mixtures were then incubated at 25°C for 10 min. The reaction was stopped with 1.0 ml of 3, 5 dinitro salicylic

acid colour reagent. The test tubes were then incubated in a boiling water bath for 5 min, cooled to room temperature. The reaction mixture was then diluted after adding 10 ml distilled water and absorbance was measured at 540 nm. Control represent 100% enzyme activity and were

conducted in similar way by replacing extract with vehicle.

Calculation of 50% inhibitory concentration (IC₅₀)

The concentration of the plant extracts required to scavenge 50% of the radicals (IC₅₀) was calculated by using the percentage scavenging activities at five different concentrations of the extract. Percentage inhibition (I %) was calculated by

$$I \% = \frac{(A_c - A_s)}{A_c} \times 100$$

where,

A_c = Absorbance of the control

A_s = Absorbance of the sample.

6.3 Thin layer chromatographic profiling

6.3.1 Practical requirements

Stationary phases

Glass plates

Mobile phase

Preparation and activation of TLC plates

Application of sample

Development tank

Development technique

Detecting or visualizing agents

6.3.1.1 Glass plates

Specific dimensions like 20 cm*20cm (full plate), 20cm*10cm (half plate), 20cm*5cm (quarter plate) can be used. It should be good quality and can withstand temperature used for drying.

6.3.1.2 Preparation and activation of TLC plates

Mixture of stationary phase and water forms slurry. TLC plates prepared by pouring, dipping, spraying or spreading. In pouring technique, slurry prepared and poured on the glass plate. Slurry spread uniformly on glass plate. Plates are dried in oven. In dipping technique two plates dipped in slurry, separated and dried. Disadvantage is that large quantity slurry required.

6.3.1.3 Spreading technique

In this technique TLC spreader is used. The glass plates are stacked on base plate. Slurry poured in the reservoir of TLC spreader. Thickness adjusted using knob in the spreader. 0.25 mm is normally used thickness for analytical purposes. Spreader rolled on the plates and air drying the plates. Plates are activated by keeping at 100 degree Celsius to 120 degree Celsius for one hour. Activated plates can be stored in thermostatically controlled oven for further use.

6.3.1.4 Application of sample

The concentration of sample of standard should be minimum for good spots. 2- 5ug of 1% solution of standard or sample spotted using capillary tubes. Spots should be 2 cm above the base of plate and spotting area should not be immersed in mobile phase. At least 4 spots can be spotted on quarter plate.

6.3.1.5 Development tank

The developing tanks require more solvent for developing tank –having hump in the middle and require less solvent. It should be lined inside with filter paper moistened with mobile phase

6.3.1.6 Mobile phase

The solvent or mobile phase depends on various factors.

Nature of substance

Nature of stationary phase

Mode of chromatography

Separation to be achieved. Analytical or preparative

6.3.2 Procedure

Commercial sheet pre coated with silica gel are available. Select a solvent by testing out the samples in various solvents. Dissolve a small quantity of ethanolic extract of *Haldina Cordifolia* leaf of the unknown in different flask containing solvents of different polarity. Place the TLC plates, the spotted side down in to the chamber so that the lower the pencil line about the solvents. Remove the plate from the development chamber and allowed to dry. Plate is placed under UV light, dark spots are observed.

6.3.3 Quantitative analysis

$$R_f = \frac{\text{Distance moved by solute}}{\text{Distance moved by solvent}}$$

R_f = retardation factor

6.4 Preliminary phytochemical screening of ethanolic leaf extract of *Haldina Cordifolia*

The ethanolic leaf extract of *Haldina Cordifolia* was used for testing preliminary phytochemical screening in order to detect major chemical groups.

6.4.1 Test for carbohydrates

- Molisch's test: Dissolved small quantity of 300mg alcoholic and dried leaf extract powder of *Pimenta dioica* separately in 4ml distilled water and filtered. The filtrate was subjected to Molisch's test.
- Fehling's test: Dissolve a small portion of extract in water and treat with Fehling's solution.
- Phenols test: The extract was spotted on a filter paper. A drop of phosphomolybdic acid reagent was added to the spot and was exposed to ammonia vapours.

6.4.2 Test for flavanoids

- Shinoda test: To 2 to 3ml of extract, a piece of magnesium ribbon and 1ml of concentrated HCl was added .
- .Lead acetate test: To 5ml of extract 1ml of lead acetate solution was added.

6.4.3Test for tannins

- Braemer's test: To a 2 to 3ml of extract, 10% alcoholic ferric chloride solution was added.

6.4.4 Test for steroid/terpenoid

- Liebermann-Burchardt test: To 1ml of extract, 1ml of chloroform, 2 to3ml of acetic anhydride and 1 to 2 drops of concentrated Sulphuric acid are added.

6.4.5 Test for alkaloids

- Draggendorf's test: A drop of extract was spotted on a small piece of precoated TLC plate and the plate was sprayed with modified Draggendorf's reagent.
- Hager's test: The extract was treated with few ml of Hager's reagen.
- Wagner's test: The extract was treated with few ml of Wagner's reagent.

6.4.6 Tests for Glycosides

- Legal's test: Dissolved the extract [0.1g] in pyridine [2ml], added sodium nitroprusside solution [2ml] and made alkaline with Sodium hydroxide solution.

6.4.7 Test for Saponins

- Foam test: 1ml of extract was dilute with 20ml of distilled water and shaken with a graduated cylinder for 15 minutes.

6.4.8 Test for Anthraquinones

- Borntrager's test: About 50 mg of powdered extract was heated with 10% ferric chloride solution and 1ml of concentrated HCl. The extract was cooled, filtered and the filtrate was shaken with diethyl ether. The ether extract was further extracted with strong ammonia.

6.4.9 Test for Amino acids

- Ninhydrin test: Dissolved a small quantity of the extract in few ml of water and added 1ml of ninhydrin reagent.

6.4.10 Test for fixed oils and fats

Press small quantity of the petroleum ether extract between two filter paper.

Note: the results for the above experiments can be noted as follows.

- If the response to the test is high it can be noted as +++which indicates that the particular group is present as the major class.
- If the response is average then note it as ++ indicates the presence in moderate quantity.
- If the response is very small then note it as + indicating the presence of only in traces.
- If no response is then negative.

6.5 Acute toxicity study

In a research study when a drug is administered to a biological system there will be some interactions may happen .In most case these are desired and usefull,but many effects are not advantageous. Acute, subacute and chronic toxicity studies are performed by the manufacturers in the investigation of a new drug.Acute toxicity is involved in estimation of LD₅₀(It is the lethal dose (cuasing death)to 50% of tested group animals)⁵⁴.

6.5.1 LD₅₀ (median lethal oral dose)

LD₅₀ (median lethal oral dose) is a statistically derived oral dose of a substance that can be expected to cause death in percent of animals when administered by the oral route. The LD₅₀ value is expressed in terms of weight of test substance per unit weight of animal (mg/kg)

In this study acute toxicity study was carried out in wistar albino rats. The procedure was followed by using OECD 423 (Acute toxic class method). The rats are fasted overnight, prior to dosing. The three dose levels are administered by the help of oral feeding needle over the prior of 24 hours. After the drugs has been administered, food may be withheld for a further 3-4 hours in rats. The purpose of sighting study is to allow selection of the appropriate starting dose for main study. The test substance is administered to a single animal in a sequential manner following from the fixed dose levels of 5, 50, 300 and 2000 mg/kg. The interval between dosing of each level is determined by the mortality/onset, duration and severity of toxic signs over the period of 24 hours, special attention given during the first 4 hours. Four hours after the drug administration, provide the food and water for 14 days and daily observed some parameters such as food intake, water intake, mortality, onset, Duration and severity of toxic signs. The animal weight is recorded on weekly once. On the day fourteen all the animals are sacrificed, to isolate the organs and observe the histopathological changes. Based on the mortality result of sighting is decided and carried out with five animals per dose level (5 or 50 or 300 or 2000 mg/kg). Based on the mortality result on 14th day of observation, the doses for invivo study are selected.

6.6 *Invivo* antidiabetic activity of *Haldina Cordifolia* leaf extract in streptozotocin induced diabetic wistar albino rats.

Wistar albino rats (150- 200 grams) of both sexes were procured from Karpagam college of pharmacy laboratory Prior to the experiment the rats were housed in a clean polypropylene cages (6 rats/ cages) for a period of 7 days under standard temperature (25 - 30⁰ c) , relative humidity (45 – 55%), dark / light cycle (12 /12 hrs). The studies were performed with the approval of Organisational Animal Ethics Committee (OAEC) (KU/IAEC/M.Pharm/173). The animals were put in overnight fasting were deprived of food for 16 hrs but allowed free access of water.

6.7 Chemicals

Streptozotocin from Loba Chemie. Standard Glibenclamide (Daonil) from Aventis Pharma. Ethanol (Analytical grade) and 5% Dextrose solution Glucose Estimation Kit from Gluco Dr Super sensor

6.8 Hypoglycemic Test

Groupings were done as follows: Group I served as control – Carboxy Methyl Cellulose (CMC) 0.5% (0.3ml\100g rat), Group II served as Positive control – Glibenclamide (2mg /kg), Group III served as aqueous ethanolic extract of *Haldina Cordifolia* – (200mg/kg), Group IV served as aqueous ethanolic extract of *Haldina Cordifolia* – (400mg/kg). Blood samples were collected by the tail nipping method and glucose level checked by glucometer. After drug Administration blood samples have been collected different time intervals at 30, 60 and 120.

6.9 Oral Glucose Tolerance Test

Groupings were done as follows: Group I served as control – Carboxy Methyl Cellulose (CMC) 0.5% (0.3ml\100g rat), Group II served as Positive control – Glibenclamide (2 mg /kg), Group III served as aqueous ethanolic extract of *Haldina Cordifolia* – (200mg/kg), Group IV served as aqueous ethanolic extract of *Haldina Cordifolia* – (400mg/kg). All the groups of animals were fasted for 24h and blood samples were collected before drug or solvent treatment. The drug, extract and solvent, have been administered to different groups and 30mins later all the groups of rats were treated with glucose orally at dose 10gm/kg body weight by using oral feeding needle. Blood samples were collected by the tail nipping method and glucose level checked by glucometer. After drug Administration blood samples have been collected different time intervals at 30, 60 and 120.

6.10 Induction of diabetes to animals

A single dose (100 mg/kg b.w., i.p.) of streptozotocin monohydrate dissolved in sodium citrate buffer was used for the induction of diabetes in rats after overnight fasting. After 1 hr of streptozotocin monohydrate administration, the animals were given feed and libitum and 5% dextrose solution was also given in feeding bottle for a day to overcome early

hypoglycaemic phase. The animals were stabilized for a week and animals showing blood glucose level more than 200 mg/dl were selected for the study.

6.11 Experimental design

Five groups of rats six in each groups received the following treatment schedule for 14 days.

- GROUP I - Normal control (normal saline 10 ml /kg, P.O)
- GROUP II - Streptozotocin treated control (100 mg/kg, I.P)
- GROUP III - Streptozotocin (100 mg/kg, I.P) +
Standard drug Glibenclamide (2 mg/kg, P.O).
- GROUP IV - Streptozotocin (100 mg/kg, i.p.) + EEVL.(200 mg/kg, P.O)
- GROUP V - Streptozotocin (100 mg/kg, i.p.)+ EEVL. (400 mg/kg, P.O)

Plant leaf extract, standard drug and normal saline were administered with the help of oral feeding needle. Group I serve as normal control which received normal saline for 14 days. Group II to Group V were diabetic control rats. Group IV and Group V (which previously received streptozotocin 100mg/kg) were given fixed doses of ethanol leaf extract (200 mg/kg, P.O, 400 mg/kg, P.O) of *Haldina Cordifolia* and group III received standard drug Glibenclamide (2 mg/kg, P.O) for 14 consecutive days. (EEKAKA-Ethanollic extract of *Haldina Cordifolia*).

6.12 Collection of blood samples

Fasting blood samples were drawn from retro orbital puncture of rats at weekly intervals till the end of the study 1, 7, and 14 days.

6.13 Estimation of biochemical parameters Serum blood glucose

On 1, 7, and 14 days fasting blood samples were collected and analyzed the blood glucose.

6.14 Blood glucose level

The blood glucose level test measures the amount of glucose in the blood sample obtained from the animals. The test is usually performed to check for elevated blood glucose levels which can be an indication of diabetes or insulin inhibition.

6.15 Statistical analysis

Statistical analysis was done by using GRAPHPAD PRISM 5.0. All the values of Biochemical parameters and body weight were expressed as Mean \pm Standard Error Mean (SEM). The values were analyzed for statistical significance using one-way analysis of variance (ANOVA), comparison was done by using Dunnett's t test. P values < 0.05 were considered as significant, P values < 0.01 were considered as very significant, P values < 0.001 were considered as highly significant and ns were considered as not significant.

CHAPTER-VII

7.RESULTS AND DISCUSSION

7.1 Appearance and percentage yield of EEHC (Ethanollic Extract of *Haldina Cordifolia* Leaf)

EEHC was a semisolid brownish colour extract and the percentage yield was found to be 14.35%

7.2 *invitro* antidiabetic study

%Inhibition of Petroleum ether extract of <i>Haldina Cordifolia</i>	leaf = 91 μ g/ml
%Inhibition of Ethyl acetate extract of <i>Haldina Cordifolia</i>	leaf = 80 μ g/ml
%Inhibition of Ethanolic extract of <i>Haldina Cordifolia</i>	leaf = 75 μ g/ml
%Inhibition of Acarbose (Positive control)	= 61 μ g/ml

Minimum % Inhibition was found in Ethanolic extract of *Haldina Cordifolia* leaf with resemblance to %Inhibition of positive control, So Ethanolic extract of *Haldina Cordifolia* contain active constituents of antidiabetic.

Table No:2, α -Amylase Inhibition of Petroleum Ether extract of *Haldina Cordifolia* leaf

Concentration(μ g/ml)	Percentage Inhibition (%)
0	0
25	29
50	33
75	42
100	53
125	56

Fig No :9




Table No:3, α -Amylase Inhibition of Ethyl acetate Extract of *Haldina Cordifolia* leaf

Concentration(μ g/ml)	Percentage Inhibition(%)
0	0
25	34
50	41
75	48
100	56
125	60

Fig No:10




Table No:4, α -Amylase Inhibition of Ethanolic Extract of *Haldina Cordifolia* leaf

Concentration(μ g/ml)	Percentage Inhibition(%)
0	0
25	30
50	36
75	50
100	56
125	61

Fig No:11





Table No:5, α -Amylase Inhibition of Acarbose (Positive control)

Concentration(μ g/ml)	Percentage Inhibition(%)
0	0
25	25
50	45
75	53
100	56
125	65

Fig No:12



7.4 TLC study

R_f Value range high-Polar substances present.

R_f value range low-Low polar substance present.

Solubility of compounds depend upon the polarity of solvents.

Obtained R_f values were confirmed by standard R_f values.

R_f values obtained for my extract ranges from 0.56 to 0.88, So my extract may contain compounds like Flavanoids, Glycosides and Alkaloids.

Table No: 6, Results of ethanolic extract of *Haldina Cordifolia*

Si No.	Solvents	Concentration	R _f value
1.	Toluene + Ethyl acetate	7:3	0.62
2.	Toluene + Ethyl acetate + Glacial acetic acid	5:5:1	0.76
3.	Petroleum ether+Chloroform	7:3	0.56
4.	Ethyl acetate + Methanol	1:1	0.88

5.	Hexane+Dichloro methane	1:1	0.67
6.	Ethyl acetate + Methanol	3:1	0.59
7.	Dichloro methane +Hexane	3:1	0.72

7.5 Phytochemical studies

The phytochemical studies results revealed that the Molisch's test no characteristic observation indicated the absence of carbohydrates, by phosphomolybdic acid test Blue coloration of the spot indicated the presence of phenols. Shinoda test and Lead acetate test gave pink or red coloration of the solution indicated the presence of flavonoids Flocculent white precipitate also indicated the same. There is no dark blue or greenish grey coloration of the solution indicated the absence of tannins in the drug. No characteristic observation for steroids and dark pink or red coloration of the solution indicated the presence of terpenoids. Orange coloration of the spot indicated the presence of alkaloids. Yellow or reddish brown precipitation precipitation indicated the presence of alkaloids. Pink to red colour solution indicates the presence of glycosides. No layer of foam formation indicates the absence of Saponins. it as + indicating the presence of traces. If no response is then negative.

Table No :7 , Results of ethanolic extract of *Haldina Cordifolia*

Class of compounds	Tests performed	Results
Carbohydrates	Molisch's test Fehling's test	-
Phenols	Phosphomolybdic acid test	+
Flavonoids	Shinoda test Lead acetate test	+ +
Tannins	Braemer's test	-
Alkaloids	Wagner's Mayer's Draggendorf's test	+ + +
Glycosides	Legal's test Brontranger's test	+ +
Saponins	Foam test	+
Sterols	Salkowski's test	-

Amino acids	Ninhydrin test	–
Terpenoids	Lieberman Burchardt test	+

+ In traces

-Absence

7.6 Hypoglycemic Test

The glucose levels were analyzed by using glucometer and each value is the mean \pm standard error (n= each group consist of 6 animals)($p<0.05$)*, ($p<0.001$)**& ($p<0.0001$)*** as compared to control & positive control group evaluated by one way, ANOVA followed by Dunnet 't' test.

The hypoglycemic test results have shown Table No:8, which indicated aqueous ethanolic extract of *Haldina Cordifolia* treated animals 200 & 400, significantly decreased in blood glucose level ($0.84 \pm 1093 \downarrow$ & $18.83 \pm 3.879 \downarrow$) ($P<0.05$)*, ($P<0.001$)**& ($P<0.0001$)*** when compared to control and positive control.

Table No-8, Hypoglycemic Test

TREATMENT	DOSE	BLOOD GLUCOSE LEVEL (mg/dl)		
		0 min	0.5hr	1 hr

	mg/kg			
CONTROL CARBOXYMETHYL CELLULOSE(CMC)	0.5 %	68.00±2.429	68.17±2.587	71.83±2.372
POSITIVE CONTROL Glibenclamide	2	69.00±0.6325	52.83±4.037**	32.83±1.515***
AQUEOUS ETHANOLIC EXTRACT OF <i>HALDINA CORDIFOLIA</i>	200	68.80±2.245	60.17±3.497*	59.33±3.584*
AQUEOUS ETHANOLIC EXTRACT OF <i>HALDINA CORDIFOLIA</i>	400	68.00±2.429	53.00±2.309**	34.17±1.138***

Fig No:13





Fig No:14



7.7 *Invivo* antidiabetic study

The experimental results have indicated on Table No:9. The negative control group glucose levels were significantly increased when compared to each other groups. All the groups of animals were affected in diabetes, which indicated blood glucose levels were slight changes in the blood glucose level ($4.13 \pm 1.207 \downarrow$ & $1. \pm 0.93 \uparrow$) for normal control group at 7th and 14th days. On day 7th glucose levels were significantly decreased glibenclamide 2mg/kg treated group ($120.2 \pm 1.414 \downarrow$ & $23 \pm 1 \downarrow$) ($P < 0.05$)*, ($P < 0.001$)** & ($P < 0.0001$)*** when compared with control group at 7th and 14th days. The ethanolic leaf extract of *Haldina Cordifolia* treated groups 200 & 400 mg/kg were dose dependent manner decreased ($P < 0.001$)** & ($P < 0.0001$)*** ($10 \pm 0.362 \downarrow$ & $90 \pm 1.67 \downarrow$) when compared with control group but positive control have more anti diabetic activity at 7th day. The aqueous ethanolic leaf extract of *Haldina Cordifolia* at the dose level 400mg/kg have equipotent activity ($90 \pm 1.67 \downarrow$ & $120.2 \pm 1.414 \downarrow$) when compared with positive control at 7th day. The ethanolic leaf extract of *Haldina Cordifolia* 200 & 400 mg/kg have been expressed dose dependent anti diabetic action ($P < 0.001$)** & ($P < 0.0001$)*** when compared to control and positive control. On day 14th, ethanolic leaf extract of *Haldina Cordifolia* treated animals 200 & 400 mg/kg significantly decreased and maintain the

blood glucose level ($5.1 \pm 0.07 \downarrow$ & $11 \pm 1.08 \downarrow$), ($P < 0.001$) ** & ($P < 0.0001$) *** when compared to control and positive control.

Table No: 9, Results of the effects of ethanolic extract on blood Glucose levels

SL. No.	TREATMENT	Blood glucose level (mg/dl) day		
		Day 1	Day 7	Day 14
1	Normal control 10 ml/kg P.O	79.83 ± 2.833	75.7 ± 4.014	76.7 ± 4.944
2	Negative control 100 mg/kg I.P	265.2 ± 3.85	270.1 ± 2.9	275.2 ± 2.5
3	Positive control (Glibenclamide 2mg/kg) P.O	255.83 ± 2.386	$135.63 \pm 3.8^{***}$	$112 \pm 2.8^{***}$
4	EEHC 200 mg/kg P.O	260 ± 3.5	$250.3 \pm 3.138^{**}$	$245.2 \pm 3.250^{**}$
5	EEHC 400 mg/kg P.O	263 ± 4.55	$173.1 \pm 2.88^{***}$	$162.1 \pm 1.8^{***}$

EEHC-Ethanolic extract of *Haldina Cordifolia* seed

(The values were expressed as Mean \pm S.E.M. (n=6 animals in each group).

7.8 Oral Glucose Tolerance Test

The glucose level were analysed by using glucometer and all values are expressed as Mean \pm SEM ($n=6$), Group 2 was compared with group 1, Groups —3,4 were compared with group 2; * $p<0.05$, ** $p<0.01$, $p<0.001$ *** evaluated by one way, ANOVA followed by Dunnet 't' test.

Oral Glucose Tolerance Test (OGTT) results have been expressed on Table No: II. Half hour after the glucose treatment, all the groups of animal blood glucose levels were significantly increased ($74.5\pm3.863\uparrow$, $35.2\pm6.6905\uparrow$, $59.5\pm3.764\uparrow$ & $65.58\pm3.762\uparrow$). The blood glucose levels were significantly decreased for , aqueous ethanolic extract of *Haldina Cordifolia* 200 & 400 mg/kg ($59.5\pm3.764\downarrow$ & $65.58\pm3.762\downarrow$) \downarrow), ($P<0.001$) **& ($P<0.0001$) ** when compared to control and positive control at 1hour and each and every $\frac{1}{2}$ hour blood glucose levels(200 mg/kg : 8.8 ± 3.26 , 10 ± 0 , 6.66 ± 1.164 & 3 ± 0.696 , 400 mg/kg: 17.9 ± 1.422 , 9.97 ± 0 , 14.7 ± 1.093 & 5 ± 0.696) ($P<0.05$)*, ($P<0.001$)**& ($P<0.0001$)*** were changes in the dose dependent manner extract treated group of animals compared to control and positive control but 400mg/kg produce the equipotent activity.

Table No-10, Oral Glucose Tolerance Test

TREATMENT	DOSE mg/kg	BLOOD GLUCOSE LEVEL (mg/dl)						
		0 min	0.5hr	1 hr	1.5hr	2 hr	2.5hr	3hr
CONTROL CARBOXY METHYL CELLULO SE(CMC)	0.5 %	68.00±2.42 ₉	142.5±6.292	187.5±9.465	172.5±12.25	157.5±12.38	153.5±12.83	130.2±13.31
POSITIVE CONTROL Glibenclam ide	2	69.00±0.63 ₂₅	104.2±7.323**	110.5±6.980***	93.67±1.308***	83.67±1.308***	77.17±4.070***	74.33±2.940***
AQUEOUS ETHANOLI C EXTRACT OF <i>HALDINA CORDIFO LIA</i>	200	68.80±2.24 ₅	128.3±6.009	147.3±2.404*	138.5±5.667*	128.5±5.667*	113.8±6.760**	108.8±6.107**
AQUEOUS METHANO LIC EXTRACT OF <i>HALDINA CORDIFO LIA</i>	400	68.00±2.42 ₉	115.0±6.191**	121.2±6.188**	103.3±4.766*** a	93.33±4.766*** a	86.67±3.602***a	83.67±2.906*** a

Fig No:15





Fig No:16



CHAPTER-VIII

8.SUMMARY AND CONCLUSION

Invitro study is on the principle of Inhibition of α -amylase, enzyme that plays a role in digestion of starch and glycogen are considered a strategy for the treatment of disorders in carbohydrate uptake, such as diabetes. Pancreatic α -amylase is a key enzyme in the digestive system and catalyses the initial step in hydrolysis of starch to a mixture of smaller. Sequential extraction was done according to increasing polarity order (Petroleum ether, Ethyl acetate and Ethanol). Each extracts were tested for α -amylase inhibition to get the extraction with minimum IC_{50} value. As per the above mechanism all the extract have concentration dependent affinity towards the inhibition of α -amylase. Finally ethanolic extract was observed as more active extract. LC-MS is a hyphenated technique, combining the separation power of HPLC, with the detection power of mass spectrometry. Even with a very sophisticated MS instrument, HPLC is still useful to remove the interferences from the sample that would impact the ionization. Closely related to LC-MS are some other techniques, like flow injection/MS, CE or CEC/MS, capillary LC or Nano LC-MS In all cases, there is the need for an interface that will eliminate the solvent and generate gas phase ions, then transferred to the optics of the mass spectrometer. Most instruments now atmospheric pressure ionization (API) technique where solvent elimination and ionization steps are combined in the source and take place at atmospheric pressure. When electron impact ionization (EI) is the choice, the solvent elimination and ionization steps are separate. Present study I have been used LC-MS for isolate and characterize the antidiabetic active compounds from extract. LC-MS for most active extract was taken and bioinformatics was collected by PASS. PASS is the computer generated program which provides the simultaneous prediction of several hundreds of biological activity types for any drug-like compounds. In this bio informatics software prediction is based on the analysis of structure-activity relationships of (SAR) the training set including more than 30000 known biologically active compounds .The most active antidiabetic constituent in the ethanolic extract of *Haldina Cordifolia* was confirmed as *Quercetin* and other antidiabetic

active constituent was confirmed as *Kaempferol*. In this present study Acute toxicity study was carried out in rats. The procedure was followed by using OECD 423 (Acute Toxic Class Method). The acute toxic class method is a step wise procedure with three animals of a single sex per step. Average two to three steps may be necessary. The method used to defined doses (2000, 1000, 500, 50, 5 mg/kg body weight, Up-and-Down Procedure). The starting dose level of EEKAGA was 2000 mg/kg body weight p.o as most of the crude extracts posses LD50 value more than 2000 mg/kg p.o and also found to be the maximum safe dose. Observe for signs for toxicity and were noted for 14 days. The onset of toxicity and signs of toxicity also noted. Hence, 1/10th (200mg/kg) and 1/5th (400mg/kg) of this dose were selected for further study. The principle involved in the alloxan induced diabetes mellitus in rats, Alloxan, a beta cytotoxic, diabetes induced chemical but wide variety of animal species by damaging the insulin secreting cells of the pancreas. Literature sources indicate that the alloxan induced rats are hyperglycaemic. The treatment of lower doses of alloxan (100mg/kg b.w.) produced partial destruction of pancreatic β -cells even though the animals become permanently diabetic. Thus these animals have surviving β -cells and regeneration is possible. It is well known that the sulfonylurea's act by directly stimulating the β -cells of the islets of Langerhans to release more insulin and these compounds are active in mild alloxan induced diabetes. *In vivo* anti diabetic screening was performed for the confirmation of above mechanism of action was undergone the ethanolic extract of *Haldina Cordifolia* in biological system (Wich was already resulted for α -amylase inhibitory activity by *In vitro* and constituents analysis by LCMS methods). At the end of the the ethanolic extract of *Haldina Cordifolia* .(200 mg/kg p.o, 400 mg/kg p.o.) showed statistically significant decrease in blood glucose levels. So the ethanolic extract of *Haldina Cordifolia* showed antidiabetic activity. This work will be useful for further diabetes mellitus and it's related diseases research worker to develop new entity for the treatment of diabetes mellitus.

Extraction of leaf was done by sequential extraction method. The leaves of *Haldina cordifolia* using the solvent with increasing polarity order (petroleum ether, ethyl acetate and ethanol) and the active extract was tested by *invitro* antidiabetic screening method. The *invitro* antidiabetic study have been performed based on the α -amylase inhibition assay. Each extracts were tested for α -amylase inhibition and the extract with minimum IC₅₀ have been undergone phyto chemical screening. The preliminary phytochemical tests was performed to identify the active phytochemicals present in the ethanolic extract of *Haldina cordifolia* and showed the presence of Phenols, Flavanoids, Alkaloids, Glycosides, Saponins and Terpenoids. LCMS analysis and PASS (bioinformatic software) was performed four compounds were found out *Kaempferol*, *Rhamnetin*, *Rhamnoxanthin*, *Quercetin* and *luteolin*. The antidiabetic constituents were found to be *Quercetin* (more active) and *Kaempferol* (less active). Finally the *invivo* antidiabetic activity of Ethanolic extract of *Haldina cordifolia* leaf was tested by using alloxan induced diabetic rat. Acute toxicity study was carried out in rats. The procedure was followed by OECD 423 (Acute toxicity class method). 1/10th (200mg/kg) and 1/5th (400mg/kg) of the maximum safe dose (2000mg/kg) were selected for further study. Fasting blood sample were drawn from retro orbital puncture of rats at weekly intervals till the end of the study 1, 7 and 14 days. On these days fasting blood glucose were collected and analysed for glucose. At the end of the study (14th day) the ethanolic extract of *Haldina cordifolia* leaf (200mg/kg p.o and 400 mg/kg p.o) treated diabetic groups showed statistically significant decrease in blood glucose similar to the standard drug glibenclamide (2mg/kg), which indicated block the α amylase activity and antagonize the alloxan action. The present study suggested that the isolation of active constituents from ethanolic extract of *Haldina cordifolia* leaf and characterize the compounds by using preliminary phytochemical studies and LCMS instrument used to isolate the compounds like *Quercetin* and *Kaempferol* were confirmed by confirmatory chemical tests.

CHAPTER-IX

9.BIBLIOGRAPHY

1. KD Tripathi MD,. 2006. Essential of medical pharmacology sixth edition.
2. American Diabetes Association Revised Fall,. 2009. Diabetes Care. Jan 2010; 33(Suppl 1): S62–S69.
3. Global health risks. Mortality and burden of disease attributable to selected major risks. Geneva, World Health Organization,2009.
4. Danaei G, Finucane MM, Lu Y, Singh GM, Cowan MJ, Paciorek CJ,. 2011. National, regional, and global trends in fasting plasma glucose and diabetes prevalence since 1980: systematic analysis of health examination surveys and epidemiological studies with 370 country-years and 2.7millionparticipants. *Lancet*,,378(9785).
5. Mathers CD, Loncar D. Projections of global mortality and burden of disease from,. 2002. to 2030. 2006, *PLoS Med*, 3(11):e442.
6. Global status report on noncommunicable diseases 2010. Geneva, World Health Organization, 2011.7. Pranav K. Prabhakar and Mukesh Doble,. 2008. A Target Based Therapeutic Approach Towards Diabetes Mellitus Using Medicinal Plants Bentham Science Publishers Ltd. *Current Diabetes Reviews*.
8. National institute of diabetes and digestive and kidney deseases ,National Diabetes Information Clearinghouse (NDIC) review 2009

9. [Shigeyuki Kawa](#), Masahiro Maruyama and Takayuki Watanabe,.2013.Prognosis and Long Term Outcomes of Autoimmune Pancreatitis ;Version 1.0, November 8,
10. Wang T.J., Larson M.G., Vasan R.S., Cheng S., Rhee E.P and McCabe E,. 2011. Metabolite profiles and the risk of developing diabetes. Nat Med.
- 11.Young K.A., Snell-Bergeon J.K., Naik R.G., Hokanson J.E., Tarullo D and Gottlieb P.A., 2011. Vitamin D deficiency and coronary artery calcification in subjects with type 1 diabetes. Diabetes Care.
12. Joergensen C., Hovind P., Schmedes A., Parving H.H and Rossing P.,2011 Vitamin d levels, microvascular complications, and mortality in type 1 diabetes. Diabetes Care.;34(5).
13. Kohei KAKU.,2010., Pathophysiology of Type 2 Diabetes and Its Treatment Policy JMAJ
14. American Diabetes Association. Standards of medical care in diabetes.,2012. *Diabetes Care*. 2012;35(Supp 1)
15. The International Expert Committee. International Expert Committee., 2009.the role of the A1C assay in the diagnosis of diabetes. Diabetes Care.
16. David b. sacks,Mark Arnold,George, Bakris,david e. bruns,Andrea rita Horvath, Kirkman and Ake Iernmark2011,Guidelines and Recommendations for Laboratory Analysis in the Diagnosis and Management of Diabetes Mellitus Diabetes Care 34:e61–e99
17. [Ronald J. Sigal](#), MD, MPH^{1,2,3},[Glen P. Kenny](#), PHD^{2,3},[David H.Wasserman](#)., 2006 .Physical Activity/Exercise and Type 2 Diabetes A consensus statement from the American Diabetes Association , Diabetes Care vol. 29 no. 6 1433-1438

18. Jane borchers.,2009. Nutrition and lifestyle recommendations for people with pre-diabetes
vol. 5 no324-45
19. Yki-JarvinenH.,2001.Combination therapies with insulin in type2 diabetes. Diabetic care.
20. Jay W. Marks, MD ., 2000. EA.New hypoglucemic therapies-J R Coll Physicians Lond.
21. Omudhome Ogbu., 2014 Insulin for Diabetes Treatment: Types, Side Effects, and PreparationsPharmacy
22. jack deRuiter 2005, Overview of the antidiabetic agents Endocrine Pharmacotherapy Module, Spring,
23. Bassam Abdul Rasool Hassan ., 2012.Medicinal Plants (Importance and Uses), 3:10
24. Hui Tag, P. Kalita, P. Dwivedi, A.K. Das and Nima D Namsa,2012. Herbal medicines used in the treatment of diabetes mellitus in Arunachal Himalaya, northeast, India [Volume 141, Issue 3](#), 14 J, Pages 786–795
25. Philip J Tusso, MD, FACP 2002, The Herbal Medicine Pharmacy Update The Permanente Journal/ Fall / Volume 6 No. 4
26. Ikuo Nishida.,2005 Journal of Plant Research ISSN: 0918-9440 ,
ISSN: 1618-0860 (electronic version)Journal.
27. Andreana L.ososki,patricia lohr,marian reiff,micheal j.balick,fredi Kronenberg and Adriane fugh-berman,bonnie o’cornor.,2002. Ethnobotanical literature survey of medicinal plants in the Dominican republic used for women’s health condition.

28. S.Dasgupta, a.parmar¹ and h.patel.,2013. International Journal of Pharma and Bio Sciences Preliminary phytochemical studies of *elaecarpus variabilis* Int J Pharm Bio Sci 4(4).

29. Isaac J. Asiedu-Gyekye , Daniel A. Antwi' Charles Awortwe , Benoit Banga N'guessan and Alexander K. Nyarko., 2007.Short-term administration of an aqueous extract of *elaecarpus variabilis*.

30. Cruz, E.A.; Reuter, S.; Martin, H.; Dehzad, N.; Muzitano, M.F.; Costa, S.S.;Rossi-Bergmann, B.; Buhl, R.; Stassen and M.; Taube.,2012 C.Phytomedicine: International Journal of Phytotherapy & Phytopharmacology.

31. Shazid M. Sharker, Mohammad K. Hossain, Mohammad R. Haque, Abu A. Chowdhury, Md. A. Kaisar, Choudhury M Hasan and Mohammad A. Rashid., 2012 Asian Pacific Journal of Tropical Biomedicine.

32. Patil SB, Dongare VR, Kulkarni CR, Joglekar MM and Arvindekar AU., 2013 Antidiabetic activity of *Elaeocarpus variabilis* in streptozotocin-induced diabetic rats by glucose independent insulin secretagogue action.

33. Gad MZ, El-Sawalhi MM, Ismail MF and El-Tanbouly ND., 2006. Biochemical study of the anti-diabetic action of the Egyptian plants fenugreek and balanites.

34. Swapnil B. Patil, Vandana R. Dongare, Chaitanya R. Kulkarni, Madhav M. Joglekar, and Akalpita U. Arvindekar.,2013. Antidiabetic activity of *Elaeocarpus variabilis* in streptozotocin-induced diabetic rats by glucose independent insulin secretagogue action.

35. [Dzhafarova RE](#), [Garaev GSh](#) and [Dzhafarkulieva ZS](#) ., 2009 . Antidiabetic action of extract of *Juglans regia*.
36. S. Radhika, Vel Sri Ranga Sanku, K.H. Smila and R. Muthezhilan., 2011. Antidiabetic and Hypolipidemic Activity of *Punica granatum* Linn on Alloxan Induced Rats World Journal of Medical Sciences.
37. Ishtiaq Ahmed, Saif-Ur-Rehman, Saghir Ahmad Jafri, Sajid Hassan, Ishtiaq Ahmed and Muhammad Naim ., 2011. Antidiabetic Effect of Aloe vera Extract on Alloxan Induced Diabetic Rats, Libyan Agriculture Research Center Journal International.
38. [Kuroda M](#), [Mimaki Y](#), [Nishiyama T](#), [Mae T](#), [Kishida H](#), [Tsukagawa M](#), [Takahashi K](#), [Kawada T](#), [Nakagawa K](#) and [Kitahara M](#) ., 2005, Hypoglycemic effects of turmeric (*Curcuma longa* L. rhizomes) on genetically diabetic KK-Ay mice.
39. Mondal, Susanta Kumar ; Chakraborty, Goutam ; Bhaumik, Uttam Kumar ; Gupta, Malaya ; Mazumder, Upal and Kanti., 2008. Antidiabetic activity of *Diospyros malabarica* Kostel bark: a preliminary investigation for possible mode of action, Oriental pharmacy and experimental medicine.
40. [Grover JK](#), Yadav.S and [Vats V J.](#), 2002. Medicinal plants of India with anti-diabetic potential, *Ethnopharmacol*.
41. Rajnish Gupta and R. S. Gupta., 2011. Antidiabetic efficacy of *mangifera indica* seed kernels in rats: a comparative study with glibenclamide, *Diabetologia Croatica*.
42. Sabjan Khaleel Basha and Vinoji Sugantha Kumari., 2012. In vitro antidiabetic activity of *psidium guajava* leaves extracts Asian Pacific Journal of Tropical Disease S98-S100.

43. Yoshino K, Miyauchi Y, Kanetaka T, Takagi Y, Koga K., 2009. Anti-diabetic activity of a leaf extract prepared from *Salacia reticulata* in mice.73(5):1096-104.
44. Akhani SP, Vishwakarma SL and Goyal RK., 2004; Anti-diabetic activity of *Zingiber officinale* in streptozotocin-induced type I diabetic rats,56(1):101-5.
45. Ramadan KS, Khalil OA, Danial EN, Alnahdi HS,and Ayaz NO., 2013.; Hypoglycemic and hepatoprotective activity of *Rosmarinus officinalis* extract in diabetic rats 69(4):779-83.
- 46.Noncommunicable Diseases in the Southeast Asia Region, Situation and Response, World Health Organization, 2011.
- 47.David R. Whiting.,2011. Global estimates of the prevalence of diabetes for 2011 and 2030, *Diabetes Research and Clinical Practice*, Volume 94, Issue 3, Pages 311-321.
48. Anjana RM, Pradeepa R, Deepa M, Datta M, Sudha V and Unnikrishnan R, 2013 “Prevalence of diabetes and prediabetes (impaired fasting glucose and/or impaired glucose tolerance) in urban and rural India. phase I results of the Indian Council of Medical Research-India Diabetes (ICMR-INDIAB) study” *Diabetologia*: a 54.12 (2011): 3022-7. NCBI.
- 49.Wunderlin,R.P and B.F..Hansen.,2008. Atlas of florida vascular plants,Institute of Systematic Botany ,FL literature *en.hortipedia.com/wiki/Elaeocarpus variabilis_elaecarpus variabilis*.
50. Michael j. Balick, Fredi kronenberg, Andreana I. Ososki,Marian reiff, Adriane fugh-berman, Bonnie o’connor,Maria roble, Patricia lohr, and Daniel atha., 2000.Medicinalal plants used by latino healers for women’shealth conditions in new medicinyork city1 *Economic Botany*.

- 51..Thalapanani N ,Chidambaram KA,Ellappan T,Sabapati ML and Mandal SC.,2008..*Journal of complimentary and Integrative medicine*.
52. Hedari R,Zareae and S,Heidarizadesh M., 2005. Journal of Nutrition.
53. Tonack S,Neoptolemos and JP,Costello E.,2010.Analysis of serum proteins by LC-MS/MS Methods, Mol Biol.
54. Glorizova,T.A Filimonov,D.A.Lagunin and A.A.Poroikov,V.,1998.Evaluation of computer system for prediction of biological activity PASS on the set of new chemical compounds.*Chim.-Pharm.J.(Rus)*.
- 55.Milne G.W.A and Miller. J.A.,1986.The information System,System overview,*JCheminf.comput.sci*.
- 56.Voigt J.H,Bienfait B,Wang S and Nicklaus M.C. .,2001.Comparison of NCI Open Database with even Large Chemical Structural Database,J.Chem.Inf Comput Sci.
57. Diabetes Blue Circle Symbol,17 March 2006. International DIABETES Federation.
- 58 .Ihlenfeldt.WD,Vogt J.H Bienfait.B,Oellien.F and Nicklaus.MC.,2002.Enhanced CACTVS Browser of the open NCI Database,j chem..inf,comput,sci.
- 59.<http://www.OECD.com>